

3. Collecting gases for pressurization

The technique used for this experiment is fairly simple: a gas is generated by adding a few mL of sulfuric acid (perhaps 5) to a small amount of solid NaNO_2 (*sample amount shown in fume hood*) in a vial. [note: this must be done in the hood and the vial should be left there after the gas has been collected]. The body of a sealed-end syringe is quickly placed over the mouth of the vial until the gas has filled it. Then it is quickly removed and stoppered with its plunger. At this point the filled syringe can be brought back to the work area. Pushing on the plunger will pressurize the gas.

What should you expect to see? If the gas is a non-mutable (doesn't change) colored material, when it is squeezed together it should look darker because more colored material is pushed into a smaller space. If there is an *equilibrium* condition with the colorless N_2O_4 then what should you expect? Will the color ever completely disappear? These are the kinds of questions you should ask yourself as you do this part of the activity. The changes in the syringe are subtle and require careful observation. Holding the syringe over white paper will help. Also, depressing the plunger quickly makes the subtle changes easier to observe. Try the process several times. [note: the changes as the gas is depressurized are generally too subtle to be observed]

The chemicals

Potassium dichromate is a bright orange-red solid, soluble in water (11.7 g per 100 g at 20°C). It is used in tanning, dyeing, painting and decorating porcelain, printing, staining wood, pyrotechnics and safety matches. The chemical is a common strong oxidizing agent for organic chemicals. It has been used medicinally as an astringent and topical antiseptic. However, it is a corrosive poison. 30 grams taken internally has been reported as fatal within 35 min. Inhalation of dust has been reported as causing cancer of the lungs.

Sodium nitrite is a white or slightly yellow crystalline solid, soluble in 1.5 parts cold water, slightly soluble in ethanol. It is decomposed by even weak acids with the evolution of brown fumes. It is used in the manufacture of dyes and in bleaching flax, silk and linen. It also had medical and veterinary applications, mainly as a relaxant for smooth muscle tissue.

Analysis

1. For reactions (1) and (3) explain your results in terms of collision theory, i.e., why do the changes in the system equilibrium position take place when you make the changes in concentration or pressure?

2. For reaction (2) explain your results with respect to a hypothetical reaction energy diagram for the reaction---sketch one and use it accompany your explanation.

3. Taken together, your observations and the generalizations that can be drawn from them constitute the main ideas behind LeChâtelier's Principle. Based on your results, generalize the effects on a system at equilibrium when the following changes are made:

- increase concentration of a reactant in a system at equilibrium
- increase concentration of a product in a system at equilibrium
- same as (a), but *decrease*
- same as (b), but *decrease*
- heat a system at equilibrium
- cool a system at equilibrium
- pressurize a gas system at equilibrium
- reduce the pressure on a gas system at equilibrium